

# **Impact of Small-Quantity Lipid-Based Nutrient Supplement on Hemoglobin, Iron Status and Biomarkers of Inflammation in Pregnant Ghanaian Women<sup>1-7</sup>**

Seth Adu-Afarwuah\*, Anna Lartey\*, Harriet Okronipa\*, Per Ashorn<sup>†</sup>, Mamane Zeilani<sup>‡</sup>, Lacey M. Baldiviez<sup>§</sup>, Brietta M. Oaks<sup>§</sup>, Stephen Vosti<sup>§</sup>, Kathryn G. Dewey<sup>§</sup>

\*Department of Nutrition and Food Science, University of Ghana, Legon, Accra, Ghana;

<sup>†</sup>University of Tampere School of Medicine, Department for International Health, Tampere,

Finland; <sup>‡</sup>Nutriset S.A.S., Hameau du Bois Ricard, PB 35, 76770 Malaunay, France; <sup>§</sup>Program in

International and Community Nutrition, Department of Nutrition, University of California,

Davis, CA, USA

<sup>1</sup> Running head: ANTENATAL LNS SUPPLEMENTATION IN GHANA.

<sup>2</sup> Funded by a grant to the University of California, Davis from the Bill & Melinda Gates Foundation. The findings and conclusions contained within are those of the authors and do not necessarily reflect positions or policies of the Bill & Melinda Gates Foundation.

<sup>3</sup> Conflict of Interest Statement: S Adu-Afarwuah, A Lartey, H Okronipa, P Ashorn, LM Baldiviez, BM Oaks, S Vosti and KG Dewey, no conflicts of interest. At the time of the study, M Zeilani was an employee of Nutriset S.A.S., which is a commercial producer of LNS products. The funder of the study had no role in the study design, data collection, analysis, and interpretation; and the preparation of the manuscript.

"This is the peer reviewed version of the following article:  
Adu-Afarwuah Seth et al. (2017) Impact of Small-Quantity Lipid-Based Nutrient Supplement on Hemoglobin, Iron Status and Biomarkers of Inflammation in Pregnant Ghanaian Women<sup>1</sup>, which has been published in final form at <http://dx.doi.org/10.1111/mcn.12262>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving."

<sup>4</sup>Address correspondence to S Adu-Afarwuah, Department of Nutrition and Food Science, University of Ghana, Legon, Accra, Ghana. Phone: +233 24 914 9385. Email: [ct3665@gmail.com](mailto:ct3665@gmail.com). Reprints are not available.

<sup>5</sup>Abbreviations: AGP, Alpha-1 glycoprotein; CRP, C-reactive protein; EFA, Essential Fatty Acid; GA, Gestational age; GW, Gestational weeks; IDA, Iron Deficiency Anemia; IFA, Iron and folic acid; ; LNS, Lipid-based Nutrient Supplement; MMN, Multiple Micronutrients; SQ-LNS, Small-quantity lipid-based nutrient supplement; TfR, Transferrin receptor; ZPP, Zinc Protoporphyrin.

<sup>6</sup>Trial Registration: ClinicalTrials.gov, Identifier NCT00970866

<sup>7</sup>Name for PubMed indexing: Adu-Afarwuah, Lartey, Okronipa, Ashorn, Zeilani, Baldiviez, Vosti, Dewey.

**Word count:** Abstract, 372; Main body of text excluding references and legends, 6,023.

### **Acknowledgements:**

We thank the families of the study subjects for their participation and cooperation; Maku Ocansey, Seth Antwi, Roney Osei-Boateing, and Boateng Bannerman for contribution to data collection and management; Janet M. Peerson for assisting with data analysis and SAS programming; Mary Arimond for comments and suggestions; iLiNS Steering Committee members Kenneth H Brown, Kenneth Maleta, and Jean Bosco Ouedraogo for advice in trial conceptualization; and Lindsay Allen for helping to define the SQ-LNS formulation and for input at various stages.

### **Contributor statement**

SA-A, AL, PA, MZ, SV, and KGD designed the research; MZ was responsible for the development and production of the LNS used in the study based on the specifications agreed

upon by the iLiNS Project; SA-A, AL, and HO conducted the research; LMB and BO performed laboratory analysis; SA-A performed the statistical analysis; AL, PA, and KGD advised on the analysis; SA-A and KGD wrote the manuscript; and AL, HO, PA, MZ, LMB, BO, and SV reviewed the draft manuscript. All authors read and approved the final manuscript.

**Abstract**

We examined hemoglobin (Hb, g/L), iron status (zinc protoporphyrin, ZPP,  $\mu\text{mol/mol}$  heme, and transferrin receptor, TfR, mg/L), and inflammation (C-reactive protein, CRP and alpha-1 glycoprotein, AGP) in pregnant Ghanaian women who participated in a randomized controlled trial. Women ( $n=1320$ ) received either 60 mg Fe + 400  $\mu\text{g}$  folic acid (IFA); 18 micronutrients including 20 mg Fe (MMN); or small-quantity lipid-based nutrient supplements (SQ-LNS, 118 kcal/d) with the same micronutrient levels as in MMN, plus 4 additional minerals (LNS) daily during pregnancy. Intention-to-treat analysis included 349, 354, and 354 women in the IFA, MMN and LNS groups, respectively, with overall baseline mean Hb and anemia (Hb  $<100$ ) prevalence of 112 and 13.3%, respectively. At 36 gestational weeks, overall Hb was 117 and anemia prevalence was 5.3%. Compared with the IFA group, the LNS and MMN groups had lower mean Hb ( $120\pm11$  vs  $115\pm12$  and  $117\pm12$ , respectively;  $p<0.001$ ), higher mean ZPP ( $42\pm30$  vs  $50\pm29$  and  $49\pm30$ ;  $p=0.010$ ) and TfR ( $4.0\pm1.3$  vs  $4.9\pm1.8$  and  $4.6\pm1.7$ ;  $p<0.001$ ), and greater prevalence of anemia (2.2% vs 7.9% and 5.8%;  $p=0.019$ ), elevated ZPP ( $>60$ ) [9.4% vs 18.6% and 19.2%;  $p=0.003$ ] and elevated TfR ( $>6.0$ ) [9.0% vs 19.2% and 15.1%;  $p=0.004$ ]. CRP and AGP concentrations did not differ among groups. We conclude that among pregnant women in a semi-urban setting in Ghana, supplementation with SQ-LNS or MMN containing 20 mg iron resulted in lower Hb and iron status but had no impact on inflammation, when compared with iron (60 mg) plus folic acid (400  $\mu\text{g}$ ). The amount of iron in such supplements that is most effective for improving both maternal Hb/iron status and birth outcomes requires further evaluation. This trial was registered at ClinicalTrials.gov as: NCT00970866.

- 23 **Keywords** lipid-based nutrient supplements, LNS, prenatal supplementation, multiple
- 24 micronutrients, hemoglobin, iron status, inflammation

## Introduction

Poor nutrient intake during pregnancy has been associated with several adverse consequences. It is estimated that up to 50% of the anemia among pregnant women in many developing country settings is due to iron deficiency (van den Broek et al., 1998) usually as a result of low dietary iron intake (World Health Organization, 1992) and poor iron bioavailability due to over-reliance on plant-based diets high in inhibitors of iron absorption such as phytate (Tatala et al., 1998). Consequences of anemia include reduced work capacity and increased risk of mortality for the mother, and premature delivery, low birth weight and poor mental development for the infant (Ren et al., 2007, International Anemia Consultative Group (INACG), 2002). While normal pregnancy is found to be associated with an increased inflammatory response (Picklesimer et al., 2008, Sacks et al., 1998), this response may be modified by macro- or micro-nutrients (Roberts et al., 2003). Higher intakes of folic acid (Bertran et al., 2005) and vitamin B<sub>6</sub> (Friso et al., 2001) have been associated with lower concentration of C-reactive protein (CRP), a common biomarker of inflammation. Several other dietary factors including essential fatty acids, EFAs (Rallidis et al., 2003) and antioxidants (Brighenti et al., 2005, Devaraj and Jialal, 2000) are also associated with the reduction of CRP concentration. Elevated CRP concentration in pregnancy is related to the development of pre-eclampsia and preterm delivery (Elovitz, 2006).

A major recommendation for increasing nutrient intake among pregnant women in developing countries is the one developed by WHO (WHO), which is the consumption of iron/folic acid (IFA) supplements containing 30-60 mg iron and 400 µg folic acid. In meta-analyses, this strategy, compared with no iron or placebo, reduced the risk of maternal anemia by 69-70 % (Imdad and Bhutta, 2012, Pena-Rosas et al., 2012) and iron deficiency by 57% (Pena-

Rosas et al., 2012), but appeared to increase the risk of reported side effects (relative risk (RR) = 2.36; 95% CI: 0.96 -5.82) particularly at iron doses of 60 mg or higher (Pena-Rosas et al., 2012). The WHO/UNICEF/UNU UNIMMAP (United Nations International Multiple Micronutrient Preparation) formulation containing 15 vitamins and minerals was more recently developed (UNICEF/WHO/UNU) to combat other possible deficiencies, e.g. for vitamins A, C and B<sub>12</sub>, which may also contribute to anemia (World Health Organization, 1992). The dose of iron in the UNIMMAP was set at 30 mg (below 60 mg) for the following reasons: (a) the presence of vitamins A, B<sub>2</sub> and C in the UNIMMAP would enhance the absorption and utilization of iron, and therefore the lower amount of iron should be sufficient, (b) a lower iron dose would be associated with less negative side effects and therefore better adherence, (c) including 60 mg of iron would mean including at least 30 mg of zinc (to avoid possible negative influence of iron on zinc absorption), bringing the total amount of metals to 90 mg, which is likely to increase negative side effects, and (d) UNIMMAP may be used in conjunction with additional iron /folic acid tablets in individual cases of more severe anemia (assuming it is caused by iron deficiency). Meta-analyses suggested that supplementation with UNIMMAP and similar products (mostly containing 11 or more micronutrients including 30 mg iron) had the same effect on maternal hemoglobin and iron status as iron (usually at 60 mg dose) with or without folic acid (Allen and Peerson, 2009, Haider et al., 2011), while also reducing the risk of low birth weight (Fall et al., 2009, Haider and Bhutta, 2012, Ramakrishnan et al., 2012).

Our group developed small quantity (20 g/d) lipid-based nutrient supplements (SQ-LNS) for pregnant and lactating women (Arimond et al., 2013) to provide micronutrients together with EFAs, using a minimum food base that supplies a small amount of energy (118 kcal/d) and high quality protein (2.6 g/d). In many populations, total energy intake among pregnant and lactating

women may be adequate, but the EFA content of the usual diet may be low (Michaelsen et al., 2011). The micronutrient composition of the SQ-LNS was generally based on the UNIMMAP formulation and a similar product used in Guinea Bissau (Kaestel et al., 2005), but we further reduced the daily iron dose to 20 mg, based on evidence that 20 mg day<sup>-1</sup> may be an adequate dose to prevent iron deficiency anemia during pregnancy (even for women who are iron deficient at entry to prenatal care) and causes fewer gastrointestinal side effects, compared to higher doses of iron (Zhou et al., 2009). We estimated (Arimond et al., 2013) that in addition to iron coming from the usual diet, the 20 mg of iron from a daily supplement would meet the recommended dietary allowance (RDA) of 27 mg iron during pregnancy (and be close to the 30 mg/d dose in the UNIMMAP formulation) while not greatly exceeding the RDA (9 mg/d) for iron during lactation (IOM, 2001, Arimond et al., 2013).

Currently, there is a growing interest in the potential use of Small-Quantity Lipid-based Nutrient Supplements (SQ-LNS) among pregnant women in developing-country settings (Hambidge et al., 2014, Research Engagement on Food Innovation for Nutritional Effectiveness (REFINE), 2013), because of evidence suggesting a positive impact of the product on certain pregnancy outcomes (Adu-Afarwuah et al., 2015). However, little is known about the impact of SQ-LNS on maternal outcomes such as anemia, iron status and inflammation. We previously reported (Adu-Afarwuah et al., 2015), that compared to IFA and a multiple micronutrient (MMN) capsule with most of the same micronutrients as the SQ-LNS, the SQ-LNS promoted fetal growth in vulnerable women, particularly primiparas, whilst the occurrence of serious adverse events did not differ between the 3 groups. In the current analysis, we compare the effect of the 3 supplementation regimens (IFA, MMN and SQ-LNS), on maternal hemoglobin (Hb),



- 93 iron status, and two biomarkers of inflammation (CRP and alpha-1 glycoprotein, AGP) during
- 94 pregnancy.

## Methods

### Study setting, design, participants and blinding

The iLiNS DYAD study in Ghana was conducted in several adjoining semi-urban communities in the Yilo Krobo and the Lower Manya Krobo Districts about 70 km north of Accra, Ghana. Details of the study setting, participants, design, randomization and masking schemes, and other key procedures have been reported elsewhere (Adu-Afarwuah et al., 2015). In brief, the study was designed as a partially double-blind, parallel, individually randomized, controlled trial with three equal-size groups. Pregnant women attending usual ante-natal clinics in four main health facilities in the area between December 2009 and December 2011 completed a screening questionnaire if they were  $\geq 18$  years old,  $\leq 20$  weeks gestation (as determined by the antenatal clinics mostly by fundal height), and had an antenatal card complete with history and examination. Informed consent for the screening was obtained by trained study workers at the antenatal clinics. Following screening, women were excluded if the antenatal card indicated HIV infection, asthma, epilepsy, tuberculosis or any malignancy. Additional exclusion criteria were known milk or peanut allergy, not residing in the area, intention to move within the next two years, unwillingness to receive field workers or take study supplement, participation in another trial, or gestational age (GA)  $> 20$  weeks before completion of the enrolment process.

Women who passed the screening were visited in their homes, where details of the study were provided, and those willing to participate were recruited, after signing or thumb-printing informed consent. Recruited women remaining eligible underwent a baseline laboratory assessment after consent, and were immediately randomized to receive one of three treatments daily: (a) 60 mg iron plus 400  $\mu$ g folic acid (hereafter, IFA supplement or group); (b) multiple micronutrient capsule containing 18 vitamins and minerals (including 20 mg iron) (hereafter,

MMN supplement or group); and (c) SQ-LNS with similar micronutrients as the MMN supplement, plus other minerals and macronutrients (hereafter, LNS supplement or group). Group allocations were developed by the Study Statistician at UC Davis using a computer-generated (SAS version 9.3) randomization scheme (in blocks of nine), and were placed in sealed, opaque envelopes. At each enrolment, a Study Nurse offered nine envelopes at a time, and the woman picked one to reveal the allocation. Allocation information was kept securely by the Field Supervisor and the Study Statistician only.

The compositions of the 3 supplements were reported previously (Adu-Afarwuah et al., 2015), as well as the considerations underlying the concentrations of the nutrients in the MMN and SQ-LNS (Arimond et al., 2013). Apart from iron which was kept at 20 mg/day in the MMN and SQ-LNS, the vitamin and mineral contents were either 1x or 2x the RDA for pregnancy, or in a few cases, the maximum amount that could be included in the supplement given technical and organoleptic constraints. The IFA and MMN supplements were provided as capsules in blister packs, and were intended to be consumed with water after a meal, one capsule per day throughout pregnancy. The LNS supplement was in 20-g sachets, and was intended to be mixed with any prepared food, one sachet per day throughout pregnancy. To maintain blinding, two individuals independent of the study placed color-coded stickers behind the blister packs (three different colors for IFA and three for MMN supplements) so that the capsules were known to the study team and participants only by the colors of the stickers. Laboratory staff and data analysts had no knowledge of group assignment until all preliminary analyses had been completed and the allocation codes were broken. The study was registered on ClinicalTrials.gov (Identifier: NCT00970866) and was approved by ethics committees of the University of California, Davis,

the Ghana Health Service, and the University of Ghana Noguchi Memorial Institute for Medical Research.

### **Procedures**

We collected socio-demographic information at baseline, and determined GA mostly by ultrasound biometry (Aloka SSD 500, Tokyo, Japan). During follow-up, field workers visited women in their homes every two weeks, whereupon they delivered a fresh supply of supplement and monitored supplement intakes. At each of laboratory assessments at baseline and at 36 GW, women's weight (Seca 874) and height (Seca 217) were measured, and peripheral malaria parasitemia (Clearview Malarial Combo, Vision Biotech, South Africa), hemoglobin, Hb (HemoCue AG, Wetzikon, Switzerland) and zinc protoporphyrin, ZPP (hematofluorometer, Aviv Biomedical Co. NJ, USA), were determined using venous blood (Adu-Afarwuah et al., 2015). We used the original Aviv cover-slides and 3-level control material for the ZPP measurements, after red blood cells were washed three times with normal saline. Plasma samples obtained after blood was centrifuged at 1,252 x g for 15 min were stored in Ghana at -20 °C, before being air-freighted on dry ice to UC Davis, where soluble transferrin receptor (TfR, mg/L), C-reactive protein (CRP, mg/L), and alpha-1 glycoprotein (AGP, g/L) concentrations were determined using a Cobas Integra 400 plus Automatic Analyzer (Roche Diagnostic Corp., Indianapolis, IN).

At 36 GW, the continuous outcomes measures were Hb (g/L), ZPP (μmol/mol heme), and plasma TfR (mg/L), CRP (mg/L), and AGP (g/L) concentrations, whilst the binary outcome measures were the percentages of women with low Hb, high Hb, and elevated ZPP, TfR, CRP and AGP.

## Sample size and data analysis

For the Ghana iLiNS-DYAD Study, an effect size (Cohen's  $d$ : difference between group means divided by the pooled standard deviation) of 0.3 (considered a small-to-moderate effect size) (Cohen) was the basis for sample size calculation. Thus, our sample size was based on detecting an effect size of 0.3 between any two groups for any continuous variable at 36 GW, with a two-sided 5% test and 80% power. As described previously (Adu-Afarwuah et al., 2015), we enrolled 1320 pregnant women into the study, but after excluding 177 who received both IFA and MMN supplements during pregnancy because of a temporary mislabeling of supplements, as well as 86 in the LNS group who were pregnant during the same time period, 1,057 women were included in the current analysis. Based on a sample size of 827 women (~275 per group) for whom data were available at 36 GW, we had 94% power to detect an effect size of 0.3 between any two groups for hemoglobin, ZPP, or TfR. This would allow a difference of 3.4 g/L in Hb, 8.9  $\mu\text{mol/mol}$  heme in ZPP, and 0.5 mg/L in TfR (given SD of 11.0, 30.0, and 2.0, respectively) to be detected between any 2 groups.

We posted the statistical analysis plan ([www.ilins.org](http://www.ilins.org)) before analysis. Statistical analysis, by intention-to-treat, was performed using SAS for Windows Release 9.3 (Cary, NC, USA). Background socio-demographic characteristics were summarized as mean  $\pm$  SD for continuous variables, or number of participants and percentages for categorical variables. As done previously (Adu-Afarwuah et al., 2015), we used 2 indices, namely assets index and housing index as proxy indicators for socioeconomic status, and calculated household food insecurity access (HFIA) score (Coates et al., 2007) as a measure of degree of household food insecurity. Higher values of the assets and housing indices represented higher socioeconomic status, and higher values of the food insecurity index represented higher food insecurity.

We calculated adherence to treatment as percentage of days from enrolment to the home visit closest to the laboratory assessment at 36 GW, when women reported consuming the supplement. We used Hb <100 g/L as our primary definition for low Hb (representing anemia). This was based on previous WHO (WHO, 2007, WHO/UNICEF/UNU, 2001) and International Nutritional Anemia Consultative Group, INACG (Nestel and INACG Steering Committee, 2002) documents that suggest lowering the standard 110 g/L cut-off by 10 g/L for pregnant women of African extraction to achieve adequate sensitivity and specificity for screening purposes (WHO/UNICEF/UNU, 2001). In addition, we defined low Hb using the standard cut-off of Hb <110 g/L, based on a recent WHO recommendation (WHO, 2011) to maintain that cut-off (110 g/L) without any adjustment, because of scarce evidence to support the adjustment. A meta-analysis (Haider et al., 2013) revealed that Hb cut-offs ranging from <100 g/L to 115 g/L have been used in studies to define anemia in pregnant women. We defined high Hb as >130 g/L (Pena-Rosas et al., 2012), elevated ZPP (proxy for iron deficiency) as >60  $\mu\text{mol/mol}$  heme (Walsh et al., 2011) and elevated TfR (proxy for tissue iron deficiency) as > 6.0 mg/L (Pfeiffer et al., 2007, Vandevijvere et al., 2013). Because there is no generally accepted cut-off value for TfR, we derived the 6.0 mg/L cut-off based on the evidence that TfR values obtained using the Automatic Analyzer assay (as used in this study) were on average 30% lower than values obtained with the ELISA assay (Pfeiffer et al., 2007). Therefore, we reduced by 30% the 8.5 mg/L cut-off value used when TfR was determined using ELISA (Vandevijvere et al., 2013) to obtain the cut-off of approximately 6.0 mg/L for our analysis. Because we used 2 cut-offs to define anemia, we also defined iron deficiency anemia (IDA) in two ways: first as Hb <100 g/L and at least one marker of iron deficiency (ZPP > 60 ( $\mu\text{mol/mol}$  heme) or TfR >6.0 mg/L (Pfeiffer et al., 2007, Vandevijvere et al., 2013)), and second, as Hb <110 g/L and at least one

marker of iron deficiency. We defined elevated inflammatory markers using the cut-off values of  $>5.0$  mg/L for CRP and  $>1.0$  g/L for AGP (Thurnham and McCabe), and categorized women with inflammation as either elevated CRP only (indicative of incubation phase of infection), elevated CRP and AGP (indicative of early convalescence) or elevated AGP only (indicative of late convalescence) (Thurnham and McCabe).

At 36 GW, we calculated overall mean ( $\pm$ SD) values and percentages for Hb and markers of iron status and inflammation. We compared groups by using general linear models (continuous outcomes) and logistic regression models (binary), with Tukey-Kramer adjustment for multiple comparisons. Along with the group comparisons, we calculated pairwise mean differences (continuous outcomes, ANOVA) and relative risks (binary outcomes, Logistic regression) with their 95% CI and p-values. Relative risks were calculated using Poisson regression (Spiegelman and Hertzmark, 2005). In addition, we analyzed changes in the prevalence of anemia, high Hb, and elevated ZPP, TfR, CRP and AGP from enrolment using mixed model logistic regression (SAS PROC GLIMMIX). Where the mixed model logistic regression failed to converge because of sparse data, we used generalized estimating equations model (SAS PROC GENMOD). We analyzed each outcome twice, first without any covariate adjustments, and then with adjustment for covariates significantly associated ( $p < 0.10$ ) with the outcome in a bivariate analysis. Because ZPP, TfR, AGP and CRP are not normally distributed, we calculated the group means ( $\pm$  SD or SE), group percentages, and pair-wise mean differences and relative risks with their 95% CI based on untransformed data, but generated the p-values for group or pair-wise comparisons using logarithmically transformed data.

To investigate the possible effect of group differences in adherence to treatment, we performed a per-protocol analysis, which was restricted to women with adherence  $\geq 70\%$ . We

evaluated potential interaction of treatment group with pre-specified baseline variables for maternal characteristics, anemia and iron status. These variables were: age, years of schooling, BMI, gestational age at enrolment, household assets index, housing index, food insecurity access score, season at enrolment (dry or wet), primiparous, anemia, and elevated ZPP, TfR, and AGP or CRP. Where an interaction was significant ( $\alpha < 0.10$ ), we performed subgroup analysis by including an interaction term between treatment and the effect modifier in the ANCOVA or logistic regression model. For continuous effect modifiers, we used data from all participants to create a linear regression model to predict the values of the outcome at the 10<sup>th</sup> and 90<sup>th</sup> percentile of the effect modifier distribution. Each effect modifier was considered separately in the models to avoid collinearity.

In a sensitivity analysis aimed at correcting for the effect of inflammation (CRP and AGP) on the Hb and iron status outcomes, we repeated the above analyses using values of Hb and iron status markers corrected for inflammation (WHO, 2007). These corrected values were calculated by grouping women into 3 inflammation categories, estimating the correction factor (CF) for each inflammation category, and multiplying the Hb and iron status values of each woman by the inflammation category-specific CF (Grant et al., 2012). The 3 inflammation categories were: reference (normal CRP and AGP), incubation (raised CRP and normal AGP), and early (raised CRP and AGP) or late (normal CRP and raised AGP) convalescence [these two phases of convalescence were combined because of small sample sizes and little indication of differences]. For ZPP at 36 GW, women were grouped into 2 inflammation categories (normal versus any inflammation), since the 3-category grouping did not yield consistent results.



## Results

We collected data from December 2009 to August 2012. The study flow diagram, as well as the main reasons women were not eligible, or were eligible but not enrolled, were reported elsewhere (Adu-Afarwuah et al., 2015). Among eligible women, those enrolled ( $n=1320$ ) and those not enrolled ( $n=606$ ) did not differ in most background characteristics (results not shown). The background characteristics of the 1057 women included in the current analysis are presented in **Table 1**. These characteristics were well-balanced across the three groups. On average, women were about 26 years of age, had about 7 years of formal education, and BMI of about 25  $\text{kg/m}^2$ . Nearly all of the women said they were married or living with a partner, slightly more than a third were primiparous, and nearly 10% tested positive for malaria. The average GA at enrolment was 16 weeks. At baseline, we obtained Hb values for all 1057 women, ZPP values for 1055 women, and TfR, CRP and AGP values for 1032 women.

At 36 GW, dropout (4.4%) was low, and did not differ among groups ( $p = 0.65$ ). Women who dropped out – mainly because of miscarriage (2.8%) and movement from the study site (1.2%) – did not differ in the baseline characteristics from those who were present at 36 GW, except for GA (weeks) at enrolment, which was significantly lower ( $p = 0.001$ ) for the former (14.5) than for the latter (16.3). Mean ( $\pm$ SD) adherence (% of days from enrolment to the home visit closest to the laboratory assessment at 36 GW when supplement was reportedly consumed) was lower ( $p=0.001$ ) in the LNS group ( $68 \pm 24$ ) compared to the IFA ( $74 \pm 21$ ) and the MMN ( $72 \pm 23$ ) groups. Women usually reported mixing the LNS supplement with porridge, but sometimes mixed it with other foods including soups and stews, or consumed it alone. We obtained Hb values for 827 women, and ZPP, TfR, CRP and AGP values for 822 women; the number (%) of women without Hb values did not differ between groups ( $p = 0.10$ ). The women

with Hb values did not differ from those without Hb values in most of the baseline characteristics, except that the latter had lower mean housing index ( $-0.17 \pm 1.09$  vs.  $0.04 \pm 0.99$ ;  $p = 0.011$ ), BMI ( $24 \pm 4.0$  vs.  $25 \pm 4.7$  kg/m<sup>2</sup>;  $p = 0.022$ ) and GA at enrolment ( $16 \pm 3.1$  vs.  $16 \pm 3.3$  weeks;  $p = 0.042$ ).

In the intention-to-treat analysis, the overall mean ( $\pm$ SD) values at baseline and 36 GW were  $112 \pm 12$  and  $117 \pm 12$ , respectively, for Hb (g/L),  $45 \pm 32$  and  $47 \pm 30$  for ZPP ( $\mu$ mol/mol heme) and  $4.1 \pm 2.5$  and  $4.5 \pm 1.7$  for TfR (mg/L). From baseline to 36 GW, the prevalence of anemia decreased significantly from 13% to 5.3% ( $p < 0.001$ ), whilst the reverse was true for the prevalence of high Hb (4.9% vs 12%,  $p < 0.001$ ) and elevated TfR (9.2% to 15%;  $p = 0.001$ ), with a moderate change in the prevalence of elevated ZPP (13% vs. 16%;  $p = 0.06$ ). The mean ( $\pm$ SD) CRP (mg/L) and AGP (g/L) at baseline ( $6.9 \pm 11$  and  $0.6 \pm 0.2$ , respectively) were slightly greater than at 36 GW ( $5.7 \pm 17$  and  $0.5 \pm 0.2$ , respectively), which was also reflected in the percentages of women with elevated CRP and AGP at baseline vs. 36 GW (38.2% vs 24.1%;  $p < 0.001$  and 6.3% vs 2.8%;  $p < 0.001$ , respectively).

## Main group comparisons at 36 gestational weeks

**Table 2** shows the unadjusted mean ( $\pm$  SD) Hb (g/L), ZPP ( $\mu$ mol/mol heme), TfR (mg/L), CRP (mg/L) and AGP (g/L) concentrations, by intervention group, at baseline and 36 GW in the intention-to-treat analysis. Baseline values did not differ significantly among groups. At 36 GW, mean Hb was significantly ( $p < 0.001$ ) greater in the IFA group ( $120 \pm 11$ ) than in the LNS ( $115 \pm 12$ ) or MMN ( $117 \pm 12$ ) group; ZPP was significantly ( $p < 0.001$ ) lower in the IFA group ( $43 \pm 30$ ) than in the LNS ( $50 \pm 29$ ) or MMN ( $49 \pm 30$ ) group; and TfR was significantly ( $p < 0.001$ ) lower in the IFA group ( $4.0 \pm 1.3$ ) than in the LNS ( $4.9 \pm 1.8$ ) or MMN ( $4.6 \pm 1.7$ ) group. Further

(**Table 3**), the percentage of women with anemia defined either as Hb <100 g/L or Hb <110 g/L was significantly lower in the IFA group compared with the LNS and MMN groups, and when using the latter definition, this percentage was also significantly greater in the LNS compared with the MMN group. Compared with the IFA group, the LNS and MMN groups had greater percentages of women with elevated ZPP (9.4% vs 19% and 19%, respectively;  $p = 0.003$ ) and elevated TfR (9.0% vs 19% and 15 %, respectively;  $p = 0.004$ ). Differences among the 3 groups in the prevalence of IDA were marginally significant ( $p = 0.07$ ) when the Hb cut-off of 100 g/L was used in the definition of IDA, but were significant ( $p < 0.001$ ) when the Hb cut-off of 110 g/L was used. In the latter situation, the risk of IDA was significantly greater in the LNS group compared with the IFA group ( $p < 0.001$ ), and marginally greater in the MMN compared to the IFA group, and in the LNS compared with the MMN group. The prevalence of high Hb did not differ among groups ( $p = 0.15$ ).

From baseline to 36 GW, the decrease in the prevalence of anemia (based on our primary definition of Hb <100 g/L) and increase in the prevalence of high Hb were significant for all groups; the change in prevalence over time differed between groups for anemia ( $P$ -interaction = 0.099) but not for high Hb ( $P$ -interaction = 0.95). For elevated ZPP, the increase in prevalence from baseline was marginally significant in the LNS group ( $p = 0.06$ ) and non-significant in the other two groups, and the change in prevalence over time did not differ between groups ( $P$ -interaction = 0.14). For elevated TfR, the increase in prevalence was significant for the LNS ( $p = 0.003$ ) and MMN ( $p = 0.009$ ) groups only, and the change in prevalence over time did not differ between groups ( $P$ -interaction = 0.17). Apart from the prevalence of anemia defined using the Hb cut-off of 110 g/L, the LNS and MMN groups did not differ significantly in any of the continuous (Table 2) or binary (Table 3) Hb and iron status outcomes.

There were no significant differences among groups in 36 GW mean ( $\pm$ SD) concentrations of CRP ( $p = 0.98$ ) and AGP ( $p = 0.35$ ) (Table 2), or the percentages of women in the incubation phase of infection (elevated CRP only;  $p = 0.26$ ), early convalescence (both CRP and AGP elevated;  $p = 0.67$ ) or late convalescence (elevated AGP only;  $p = 1.00$ ). From baseline to 36 GW, the decrease in the percentage of women in the incubation phase of infection was significant for all groups. For percentage of women in early convalescence, only the decrease in the IFA group ( $p = 0.010$ ) was significant, and for percentage of women in late convalescence (where the mixed model logistic regression did not converge because of sparse data, and hence generalized estimating equations model was used), only the decrease in the MMN group ( $p = 0.024$ ) was significant.

Adjustments by covariates significantly associated with Hb and the iron status (ZPP and TfR) and inflammatory (CRP and AGP) outcomes (including the baseline value for each outcome) did not alter the unadjusted results (data not shown). In addition, correcting the Hb and iron status values for inflammation in the sensitivity analysis (results not shown) did not change the above findings.

The per-protocol analysis (results not shown) revealed that among women with adherence to treatment  $\geq 70\%$  (samples sizes at 36 gestational weeks: 213 in IFA, 222 in MMN and 166 in the LNS group, for Hb), the above findings remained unchanged, except for the fact that the prevalence of anemia (defined as Hb  $< 100$  g/L or  $< 110$  g/L) did not differ between the LNS and MMN groups, and the prevalence of IDA (if definition included Hb  $< 110$  g/L) was greater only in the LNS compared with the IFA group.

#### **Effect modification**

Interactions of treatment group with BMI, season of enrolment and elevated CRP

concentration at baseline were not significant for any of the outcomes. As shown in **Table 4**, the effect of intervention on ZPP concentration at 36 GW was modified by baseline anemia ( $Hb < 100$  g/L;  $P$ -interaction = 0.099), elevated ZPP ( $P$ -interaction = 0.013) and TfR ( $P$ -interaction = 0.090), GA at enrolment ( $P$ -interaction = 0.041), and household assets score ( $P$ -interaction = 0.061). Specifically, the difference in mean ( $\pm$ SD) ZPP concentration at 36 GW among intervention groups (MMN and LNS groups compared with the IFA group) was greater among women with anemia at baseline, elevated baseline ZPP, greater GA at enrolment, or lower household assets scores. Similarly, the difference in risk of elevated ZPP among intervention groups was greater in women with greater GA at enrolment ( $P$ -interaction = 0.057), and elevated TfR at baseline ( $P$ -interaction = 0.049)

The difference in mean TfR concentration at 36 GW among intervention groups was greater among women who were anemic at baseline ( $P$ -interaction = 0.079), did not have elevated AGP at baseline ( $P$ -interaction = 0.051), or had greater GA at enrolment ( $P$ -interaction = 0.016), whereas the difference in risk of elevated TfR among intervention groups was greater in women with less food insecurity.

For elevated CRP and AGP at 36 GW, there were significant interactions between intervention group and elevated ZPP at baseline ( $P$ -interaction = 0.008) and household assets score ( $P$ -interaction = 0.091), but the stratified analyses (Table 4) did not show consistent results.

## Discussion

In the iLiNS-DYAD-Ghana study, pregnant women who were provided with standard iron (60 mg) and folic acid supplements from  $\leq 20$  GW had significantly greater mean Hb, lower mean ZPP and TfR and lower prevalence of anemia and iron deficiency (elevated ZPP and TfR) at 36 GW than pregnant women who were provided with either the MMN or LNS supplements, both of which contained 20 mg iron. Overall, however, the prevalence of anemia (Hb < 100 g/L) at 36 GW was relatively low (2.2-7.9%), and iron deficiency was evident in <20% of women in all groups. At 36 GW, the three groups did not differ in the percentage of women with high Hb (12.3% overall) or inflammation (CRP and AGP). These findings remained unchanged when analyses were restricted to women who were more adherent to treatment.

A few weaknesses of our study were described previously (Adu-Afarwuah et al., 2015), namely: (a) a fully double-blind study design was not possible because of the physical differences between the SQ-LNS in the form of sachets, and the other two supplements (MMN and IFA) in the form of capsules, and (b) adherence to treatment was assessed by self-report and not direct observation. However, none of the individuals involved in sample collection, laboratory measurements or data analysis had any knowledge of group assignment. The good collaboration with the antenatal clinics (which gave women the confidence to cooperate), relatively low rate of attrition, intense follow-up of participants, and detailed attention we paid to ensuring data quality were notable strengths of the study.

Several possible explanations may be relevant for the observed lower Hb and higher ZPP and TfR values in the MMN and LNS groups compared to the IFA group. First, the 20 mg iron dose used in the MMN and LNS supplements may have been too low for this Ghanaian population of pregnant women. In most (but not all) similar studies, the iron dose of the iron +

folic acid supplement was 60 mg/day and that of the multiple micronutrient supplement was 30 mg/day (Allen and Pearson, 2009, Roberfroid et al., 2011, Mei et al., 2014), although in Indonesia (Suprpto et al., 2002), Mexico (Ramakrishnan et al., 2004), Nepal (Christian et al., 2003) and Tanzania (Makola et al., 2003), both the iron + folic acid and multiple micronutrient supplement groups received iron doses of at least 50 mg/ day. In these studies (even for those that used multiple micronutrients containing 30 mg Fe), pregnant women consuming the multiple micronutrient supplements generally did not differ in Hb or iron status indicators compared to those consuming iron + folic acid (Allen and Pearson, 2009, Mei et al., 2014, Roberfroid et al., 2011). A study of Australian women (Zhou et al., 2009) suggested that 20 mg iron per day may be an adequate dose to prevent iron deficiency anemia during pregnancy compared with higher doses of iron. However, the diet of the women in our sample, as is typical of Ghana, is mainly plant-based and high in phytate (Gibson, 1994), which reduces iron absorption (Baech et al., 2003), so dietary iron needs during pregnancy may be higher in Ghana than in Australia.

Another possibility is that the relatively high dose of zinc (30 mg) in the MMN and LNS supplements may have interfered with iron absorption, as suggested by the results of a supplementation trial in Nepal (Christian et al., 2003). The lack of a good biomarker of zinc status at the individual level makes it difficult to explore this potential mechanism for the differences in iron status between the IFA and other two groups.

It is noteworthy that there were no significant differences in mean Hb or iron status between the LNS and MMN groups, despite the fact that SQ-LNS is a food-based supplement rather than a capsule and some differences in composition could have affected these outcomes. For instance, it is possible that the calcium or phytate in SQ-LNS could have limited the absorption of iron in the LNS group. The lack of differences in Hb and iron status between these

two groups suggests that the iron or other micronutrient content (which was identical in these two supplements, except for the macro-minerals) was the most critical factor.

In Ghana, anemia (Ghana Statistical Service (GSS) et al., 2009) and infections including malaria (Yatich et al., 2009) are common among pregnant women even in relatively high income communities, and evidence (Mockenhaupt et al., 2000) suggests that malaria is a major risk factor for anemia. Thus, it is noteworthy that the prevalence of both anemia and elevated CRP declined significantly in all three groups between baseline and 36 weeks gestation.

It is important to consider the implications of the observed lower mean Hb of the SQ-LNS and MMN groups compared to the IFA group at 36 weeks gestation. Low Hb concentration or anemia in the first or second trimester of pregnancy is linked with poor pregnancy outcomes including low birth weight (Murphy et al., 1986), but no such association has been established for low Hb concentration or anemia in the third trimester (Allen, 2000). Further, Hb concentrations substantially above 110–119 g/L during pregnancy may be independent of iron status and have been linked with poorer health outcomes for the mother and fetus (Yip, 2000, Zhou et al., 1998). Therefore, the higher mean Hb concentrations observed for women in the IFA group compared to those in the LNS and MMN groups in the third trimester (36 GW) may not necessarily be beneficial with respect to birth outcomes. In fact, we previously demonstrated that in this same study (Adu-Afarwuah et al., 2015) the prenatal consumption of LNS (compared to IFA) was associated with greater birth weight, weight-for-age z-score and BMI-for-age z-score, and that, in first-time mothers, prenatal LNS supplementation also increased birth length and head circumference and reduced the proportion of infants with low birth weight, low birth length, and small-for-gestational age. In this cohort, there was no relationship between Hb at 36 wk gestation and infant birth size, and there was actually a significant *negative* relationship between



maternal iron status at 36 wk and birth size (Oaks et al., 2015). Thus, the difference between the LNS and IFA groups in mean maternal Hb, ZPP, and TfR concentrations needs to be weighed against the difference (in the opposite direction) in birth outcomes. In two sets of analyses (Garn et al., 1981, Steer et al., 1995) each involving a large number of pregnant women, the lowest risk of adverse birth outcomes including low birth weight was seen in women with Hb ~95-105 g/L (Steer et al., 1995) or Hb ~100 -110 g/L (Garn et al., 1981). Appropriate cut-offs for ZPP and TfR in pregnancy are not well documented, particularly with respect to functional outcomes. Therefore, it is difficult to judge whether our results should be interpreted as “improvements” in maternal iron status in the IFA group compared to the MMN or LNS group.

We conclude that among pregnant women in a semi-urban setting in Ghana, supplementation with SQ-LNS or MMN containing 20 mg iron resulted in lower Hb and iron status but had no impact on inflammation, when compared with iron (60 mg) plus folic acid (400 µg) treatment. The amount of iron in such supplements that is most effective for improving both maternal Hb/iron status and birth outcomes requires further evaluation.

#### **Key messages**

1. In this semi-urban Ghanaian population, the prevalence of anemia (Hb <100 g/L) among pregnant women who received IFA, MMN or SQ-LNS was reduced from 13% at <20 gestational weeks (GW) to 5% at 36 GW.
2. Provision of IFA (with 60 mg Fe) was associated with a greater mean concentration of Hb and lower prevalence of anemia at 36 GW than provision of MMN or SQ-LNS (both with 20 mg Fe).

- 457 3. The prevalence of high Hb (>130 g/L) or elevated inflammatory biomarkers (CRP and  
458 AGP) at 36 GW was not affected by the type of prenatal supplement provided.
- 459 4. More research is needed to determine the concentration of iron in MMN and SQ-LNS  
460 supplements that is most effective for improving both maternal Hb/iron status and birth  
461 outcomes.

**References**

- Adu-Afarwuah S., Lartey A., Okronipa H., Ashorn P., Zeilani M., Pearson J.M., et al. (2015) Lipid-based nutrient supplement increases the birth size of infants of primiparous women in Ghana. *The American Journal of Clinical Nutrition* **101**, 835-846.
- Allen L.H. (2000) Anemia and iron deficiency: effects on pregnancy outcome. *The American Journal of Clinical Nutrition* **71**, 1280S-1284S.
- Allen L.H. & Pearson J.M. (2009) Impact of multiple micronutrient versus iron-folic acid supplements on maternal anemia and micronutrient status in pregnancy. *Food and Nutrition Bulletin* **30**, S527-532.
- Arimond M., Zeilani M., Jungjohann S., Brown K.H., Ashorn P., Allen L.H., et al. (2013) Considerations in developing lipid-based nutrient supplements for prevention of undernutrition: experience from the International Lipid-Based Nutrient Supplements (iLiNS) Project. *Maternal & Child Nutrition*.
- Baech S.B., Hansen M., Bukhave K., Jensen M., Sorensen S.S., Kristensen L., et al. (2003) Nonheme-iron absorption from a phytate-rich meal is increased by the addition of small amounts of pork meat. *The American Journal of Clinical Nutrition* **77**, 173-179.
- Bertran N., Camps J., Fernandez-Ballart J., Arija V., Ferre N., Tous M., et al. (2005) Diet and lifestyle are associated with serum C-reactive protein concentrations in a population-based study. *Journal of Laboratory and Clinical Medicine* **145**, 41-46.
- Brighenti F., Valtuena S., Pellegrini N., Ardigo D., Del Rio D., Salvatore S., et al. (2005) Total antioxidant capacity of the diet is inversely and independently related to plasma concentration of high-sensitivity C-reactive protein in adult Italian subjects. *The British Journal of Nutrition* **93**, 619-625.

- 485 Christian P., Shrestha J., LeClerq S.C., Khatry S.K., Jiang T., Wagner T., et al. (2003)  
486       Supplementation with micronutrients in addition to iron and folic acid does not further  
487       improve the hematologic status of pregnant women in rural Nepal. *The Journal of*  
488       *Nutrition* **133**, 3492-3498.
- 489 Coates J., Swindale A. & Bilinsky P. (2007) Household Food Insecurity Access Scale (HFIAS)  
490       for Measurement of Food Access: Indicator Guide (V.3) [Internet]. Food and Nutrition  
491       Technical Assistance Project, Academy for Educational Development, Washington, D.C  
492       [cited 2013 Aug 12]. Available from: [http://www.fao.org/fileadmin/user\\_upload/eufao-](http://www.fao.org/fileadmin/user_upload/eufao-fsi4dm/doc-training/hfias.pdf)  
493       [fsi4dm/doc-training/hfias.pdf](http://www.fao.org/fileadmin/user_upload/eufao-fsi4dm/doc-training/hfias.pdf).
- 494 Cohen J. Statistical Power Analysis in the Behavioral Sciences (2nd edition). Hillsdale (NJ):  
495       Lawrence Erlbaum Associates, Inc., 1988.
- 496 Devaraj S. & Jialal I. (2000) Alpha tocopherol supplementation decreases serum C-reactive  
497       protein and monocyte interleukin-6 levels in normal volunteers and type 2 diabetic  
498       patients. *Journal of Free Radicals in Biology & Medicine* **29**, 790-792.
- 499 Elovitz M.A. (2006) Anti-inflammatory interventions in pregnancy: Now and the future.  
500       *Seminars in Fetal & Neonatal Medicine* **11**, 327-332.
- 501 Fall C.H., Fisher D.J., Osmond C. & Margetts B.M. (2009) Multiple micronutrient  
502       supplementation during pregnancy in low-income countries: a meta-analysis of effects on  
503       birth size and length of gestation. *Food and Nutrition Bulletin* **30**, S533-546.
- 504 Friso S., Jacques P.F., Wilson P.W.F., Rosenberg I.H. & Selhub J. (2001) Low Circulating  
505       Vitamin B6 Is Associated With Elevation of the Inflammation Marker C-Reactive Protein  
506       Independently of Plasma Homocysteine Levels. *Circulation* **103**, 2788-2791.

- 507 Garn S.M., Keating M.T. & Falkner F. (1981) Hematological status and pregnancy outcomes.  
508 *The American Journal of Clinical Nutrition* **34**, 115-117.
- 509 Ghana Statistical Service (GSS), Ghana Health Service (GHS) & ICF Macro (2009) Ghana  
510 Demographic and Health Survey 2008. Accra, Ghana: GSS, GHS, and ICF Macro.
- 511 Gibson R.S. (1994) Zinc nutrition in developing countries. *Nutrition Research Reviews* **7**, 151-  
512 173.
- 513 Grant F.K., Suchdev P.S., Flores-Ayala R., Cole C.R., Ramakrishnan U., Ruth L.J., et al. (2012)  
514 Correcting for inflammation changes estimates of iron deficiency among rural Kenyan  
515 preschool children. *The Journal of Nutrition* **142**, 105-111.
- 516 Haider B.A. & Bhutta Z.A. (2012) Multiple-micronutrient supplementation for women during  
517 pregnancy. *The Cochrane Database of Systematic Reviews* **11**, CD004905.
- 518 Haider B.A., Olofin I., Wang M., Spiegelman D., Ezzati M., Fawzi W.W., et al. (2013) Anaemia,  
519 prenatal iron use, and risk of adverse pregnancy outcomes: systematic review and meta-  
520 analysis. *BMJ*, **346**, f3443.
- 521 Haider B.A., Yakoob M.Y. & Bhutta Z.A. (2011) Effect of multiple micronutrient  
522 supplementation during pregnancy on maternal and birth outcomes. *BMC Public Health*  
523 **11 Suppl 3**, S19.
- 524 Hambidge K.M., Krebs N.F., Westcott J.E., Garces A., Goudar S.S., Kodkany B.S., et al. (2014)  
525 Preconception maternal nutrition: a multi-site randomized controlled trial. *BMC*  
526 *Pregnancy & Childbirth* **14**, 111.
- 527 Imdad A. & Bhutta Z.A. (2012) Routine iron/folate supplementation during pregnancy: effect on  
528 maternal anaemia and birth outcomes. *Paediatric and Perinatal Epidemiology* **26 Suppl**  
529 **1**, 168-177.

- 530 International Anemia Consultative Group (INACG) (2002) Report of the 2001 International  
531 Anemia Consultative Group Symposium. Why is iron important and what to do about it:  
532 a new perspective. Washington, DC, INACG Secretariat.
- 533 IOM (2001) *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium,*  
534 *Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium and Zinc.*  
535 National Academy Press, Washington, DC.
- 536 Kaestel P., Michaelsen K.F., Aaby P. & Friis H. (2005) Effects of prenatal multimicronutrient  
537 supplements on birth weight and perinatal mortality: a randomised, controlled trial in  
538 Guinea-Bissau. *The European Journal of Clinical Nutrition* **59**, 1081-1089.
- 539 Makola D., Ash D.M., Tatala S.R., Latham M.C., Ndossi G. & Mehansho H. (2003) A  
540 micronutrient-fortified beverage prevents iron deficiency, reduces anemia and improves  
541 the hemoglobin concentration of pregnant Tanzanian women. *The Journal of Nutrition*  
542 **133**, 1339-1346.
- 543 Mei Z., Serdula M.K., Liu J.M., Flores-Ayala R.C., Wang L., Ye R., et al. (2014) Iron-  
544 containing micronutrient supplementation of Chinese women with no or mild anemia  
545 during pregnancy improved iron status but did not affect perinatal anemia. *The Journal of*  
546 *Nutrition* **144**, 943-948.
- 547 Michaelsen K.F., Dewey K.G., Perez-Exposito A.B., Nurhasan M., Lauritzen L. & Roos N.  
548 (2011) Food sources and intake of n-6 and n-3 fatty acids in low-income countries with  
549 emphasis on infants, young children (6-24 months), and pregnant and lactating women.  
550 *Maternal & Child Nutrition* **7 Suppl 2**, 124-140.
- 551 Mockenhaupt F.P., Rong B., Günther M., Beck S., Till H., Kohne E., et al. (2000) Anaemia in  
552 pregnant Ghanaian women: importance of malaria, iron deficiency, and

- 553 haemoglobinopathies. *Transaction of the Royal Society of Tropical Medicine & Hygiene*  
554 **94**, 477-483.
- 555 Murphy J.F., O'Riordan J., Newcombe R.G., Coles E.C. & Pearson J.F. (1986) Relation of  
556 haemoglobin levels in first and second trimesters to outcome of pregnancy. *Lancet* **1**,  
557 992-995.
- 558 Nestel P. & INACG Steering Committee (2002) Adjusting Hemoglobin Values in Program  
559 Surveys [Internet]. INACG, Washington, DC [cited 2014-Jan 10]. Available from:  
560 [http://pdf.usaid.gov/pdf\\_docs/PNACQ927.pdf](http://pdf.usaid.gov/pdf_docs/PNACQ927.pdf).
- 561 Oaks B., Stewart C., Laugero K., Adu-Afarwuah S., Lartey A., Baldiviez L., et al. (2015)  
562 Associations of maternal cortisol, inflammation, hemoglobin, iron status, and BMI with  
563 birth outcomes in pregnant women in Ghana. *The Federation of American Societies for*  
564 *Experimental Biology Journal* **29**.
- 565 Pena-Rosas J.P., De-Regil L.M., Dowswell T. & Viteri F.E. (2012) Daily oral iron  
566 supplementation during pregnancy. *The Cochrane Database of Systematic Reviews* **12**,  
567 CD004736.
- 568 Pfeiffer C.M., Cook J.D., Mei Z., Cogswell M.E., Looker A.C. & Lacher D.A. (2007) Evaluation  
569 of an automated soluble transferrin receptor (sTfR) assay on the Roche Hitachi analyzer  
570 and its comparison to two ELISA assays. *Clinica Chimica Acta; International Journal of*  
571 *Clinical Chemistry* **382**, 112-116.
- 572 Picklesimer A.H., Jared H.L., Moss K., Offenbacher S., Beck J.D. & Boggess K.A. (2008)  
573 Racial differences in C-reactive protein levels during normal pregnancy. *American*  
574 *Journal of Obstetrics and Gynecology* **199**, 523.e521-523.e526.

- 575 Rallidis L.S., Paschos G., Liakos G.K., Velissaridou A.H., Anastasiadis G. & Zampelas A.  
576 (2003) Dietary  $\alpha$ -linolenic acid decreases C-reactive protein, serum amyloid A and  
577 interleukin-6 in dyslipidaemic patients. *Atherosclerosis* **167**, 237-242.
- 578 Ramakrishnan U., Grant F.K., Goldenberg T., Bui V., Imdad A. & Bhutta Z.A. (2012) Effect of  
579 multiple micronutrient supplementation on pregnancy and infant outcomes: a systematic  
580 review. *Paediatric and Perinatal Epidemiology* **26 Suppl 1**, 153-167.
- 581 Ramakrishnan U., Neufeld L.M., Gonzalez-Cossio T., Villalpando S., Garcia-Guerra A., Rivera  
582 J., et al. (2004) Multiple micronutrient supplements during pregnancy do not reduce  
583 anemia or improve iron status compared to iron-only supplements in Semirural Mexico.  
584 *The Journal of Nutrition* **134**, 898-903.
- 585 Ren A., Wang J., Ye R.W., Li S., Liu J.M. & Li Z. (2007) Low first-trimester hemoglobin and  
586 low birth weight, preterm birth and small for gestational age newborns. *International*  
587 *Journal of Gynaecology and Obstetrics: The Official Organ of the International*  
588 *Federation of Gynaecology and Obstetrics* **98**, 124-128.
- 589 Research Engagement on Food Innovation for Nutritional Effectiveness (REFINE) (2013)  
590 Ongoing studies using Lipid Based Nutrient Supplement (LNS) [Internet]. REFINE.  
591 Boston, MA. [cited 2015 Jun 23]. Available from:  
592 <http://refinenutrition.org/research/REFINE-product-LNS.pdf>.
- 593 Roberfroid D., Huybregts L., Habicht J.P., Lanou H., Henry M.C., Meda N., et al. (2011)  
594 Randomized controlled trial of 2 prenatal iron supplements: is there a dose-response  
595 relation with maternal hemoglobin? *The American Journal of Clinical Nutrition* **93**,  
596 1012-1018.



- 597 Roberts J.M., Balk J.L., Bodnar L.M., Belizán J.M., Bergel E. & Martinez A. (2003) Nutrient  
598 Involvement in Preeclampsia. *The Journal of Nutrition* **133**, 1684S-1692S.
- 599 Sacks G.P., Studena K., Sargent I.L. & Redman C.W.G. (1998) Normal pregnancy and  
600 preeclampsia both produce inflammatory changes in peripheral blood leukocytes akin to  
601 those of sepsis. *American Journal of Obstetrics and Gynecology* **179**, 80-86.
- 602 Spiegelman D. & Hertzmark E. (2005) Easy SAS calculations for risk or prevalence ratios and  
603 differences. *American Journal of Epidemiology* **162**, 199-200.
- 604 Steer P., Alam M.A., Wadsworth J. & Welch A. (1995) Relation between maternal haemoglobin  
605 concentration and birth weight in different ethnic groups. *BMJ*, **310**, 489-491.
- 606 Suprpto B., Widardo & Suhanantyo (2002) Effect of low-dosage vitamin A and riboflavin on  
607 iron-folate supplementation in anaemic pregnant women. *Asia Pacific Journal Clinical*  
608 *Nutrition* **11**, 263-267.
- 609 Tatala S., Svanberg U. & Mduma B. (1998) Low dietary iron availability is a major cause of  
610 anemia: a nutrition survey in the Lindi District of Tanzania. *The American Journal of*  
611 *Clinical Nutrition* **68**, 171-178.
- 612 Thurnham D.I. & McCabe G.P. Influence of infection and inflammation on biomarkers of  
613 nutritional status with an emphasis on vitamin A and iron. In: World Health  
614 Organization. Report: Priorities in the assessment of vitamin A and iron status in  
615 populations, Panama City, Panama, 15–17 September 2010 [Internet]. World Health  
616 Organization, Geneva; 2012 [cited 2014 Nov 14]. Available from:  
617 [http://www.who.int/nutrition/publications/micronutrients/background\\_paper4\\_report\\_assessment\\_vitAandIron\\_status.pdf](http://www.who.int/nutrition/publications/micronutrients/background_paper4_report_assessment_vitAandIron_status.pdf).  
618

- 619 UNICEF/WHO/UNU Composition of a multi-micronutrient supplement to be used in pilot  
620 programmes among pregnant women in developing countries [Internet]. New York:  
621 UNICEF. 1999 [cited 2015 Jan 18]. Available from:  
622 [http://apps.who.int/iris/bitstream/10665/75358/1/UNICEF-WHO-multi-](http://apps.who.int/iris/bitstream/10665/75358/1/UNICEF-WHO-multi-micronutrients.pdf?ua=1)  
623 [micronutrients.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/75358/1/UNICEF-WHO-multi-micronutrients.pdf?ua=1).
- 624 van den Broek N.R., Letsky E.A., White S.A. & Shenkin A. (1998) Iron status in pregnant  
625 women: which measurements are valid? *The British Journal of Haematology* **103**, 817-  
626 824.
- 627 Vandevijvere S., Amsalkhir S., Van Oyen H., Egli I. & Moreno-Reyes R. (2013) Iron status and  
628 its determinants in a nationally representative sample of pregnant women. *Journal of the*  
629 *Academy of Nutrition and Dietetics* **113**, 659-666.
- 630 Walsh T., O'Broin S.D., Cooley S., Donnelly J., Kennedy J., Harrison R.F., et al. (2011)  
631 Laboratory assessment of iron status in pregnancy. *Clinical Chemistry and Laboratory*  
632 *Medicine : CCLM / FESCC* **49**, 1225-1230.
- 633 WHO Guideline: Daily iron and folic acid supplementation in pregnant women [Internet]. World  
634 Health Organization, Geneva; 2012 [cited 2014 Jul 13]. Available from:  
635 [http://apps.who.int/iris/bitstream/10665/77770/1/9789241501996\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/77770/1/9789241501996_eng.pdf?ua=1).
- 636 WHO (2007) Assessing the iron status of populations: Report of a joint World Health  
637 Organization/ Centers for Disease Control and Prevention technical consultation on the  
638 assessment of iron status at the population level, 2nd Ed. [Internet]. World Health  
639 Organization, Geneva [cited 2015 Oct 22]. Available from:  
640 [http://www.who.int/nutrition/publications/micronutrients/anaemia\\_iron\\_deficiency/97892](http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/9789241596107.pdf)  
641 [41596107.pdf](http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/9789241596107.pdf)

- WHO (2011) Hemoglobin concentrations for the diagnosis of anemia and assessment of severity [Internet]. Vitamin and Mineral Nutrition Information System. Geneva, World Health Organization (WHO/NMH/NHD/MNM/11.1) [cited 2015 Oct 21]. Available from: <http://www.who.int/vmnis/indicators/haemoglobin.pdf>.
- WHO/UNICEF/UNU (2001) Iron deficiency anaemia assessment, prevention, and control: a guide for programme managers. WHO/NHD/01.3. [Internet]. World Health Organization, Geneva [cited 2014 Dec 01]. Available from: [http://www.who.int/nutrition/publications/en/ida\\_assessment\\_prevention\\_control.pdf](http://www.who.int/nutrition/publications/en/ida_assessment_prevention_control.pdf).
- World Health Organization (1992) The Prevalence of Anaemia in Women. WHO/MCH /MSM/92.2.
- Yatich N.J., Yi J., Agbenyega T., Turpin A., Rayner J.C., Stiles J.K., et al. (2009) Malaria and Intestinal Helminth Co-infection Among Pregnant Women in Ghana: Prevalence and Risk Factors. *The American Journal of Tropical Medicine and Hygiene* **80**, 896-901.
- Yip R. (2000) Significance of an abnormally low or high hemoglobin concentration during pregnancy: special consideration of iron nutrition. *The American Journal of Clinical Nutrition* **72**, 272S-279S.
- Zhou L.-M., Yang W.-W., Hua J.-Z., Deng C.-Q., Tao X. & Stoltzfus R.J. (1998) Relation of hemoglobin measured at different times in pregnancy to preterm birth and low birth weight in Shanghai, China. *American Journal of Epidemiology* **148**, 998-1006.
- Zhou S.J., Gibson R.A., Crowther C.A. & Makrides M. (2009) Should we lower the dose of iron when treating anaemia in pregnancy? A randomized dose-response trial. *European Journal of Clinical Nutrition* **63**, 183-190

**TABLE 1**

Background characteristics of pregnant Ghanaian women whose hemoglobin, and iron status and inflammatory markers were analyzed at 36 gestational weeks, by intervention group<sup>1</sup>

Background characteristics	IFA (N = 349)	MMN (N = 354)	LNS (N = 354)
Age, y	27 ± 5.3 [349]	27 ± 5.7 [354]	27 ± 5.4 [354]
Formal education, y	7.6 ± 3.5 [349]	7.5 ± 3.6 [354]	7.7 ± 3.7 [354]
Body Mass Index, kg/m <sup>2</sup>	25 ± 4.3 [342]	25 ± 5.0 [348]	25 ± 4.4 [349]
Low BMI, n/N (%)	11/342 (3.2)	8/348 (2.3)	6/349 (1.7)
Gestational age at enrolment, weeks	16.3 ± 3.3 [346]	16.2 ± 3.2 [353]	16.2 ± 3.3 [349]
Assets index <sup>2</sup>	0.09 ± 0.98 [342]	0.1 ± 0.9 [349]	-0.01 ± 0.91 [348]
Housing index <sup>2</sup>	0.04 ± 0.99 [342]	-0.03 ± 1.03 [349]	0.00 ± 1.00 [348]
HFIA Score <sup>3</sup>	2.9 ± 4.6 [345]	2.5 ± 3.9 [346]	2.5 ± 3.9 [348]
Married or co-habiting, n/N (%)	320/349 (91.7)	332/354 (93.8)	328/354 (92.7)
Primiparous women, n/N (%)	131/349 (37.5)	110/354 (31.1)	128/354 (36.2)
Tested positive for malaria <sup>4</sup> , n/N (%)	31/349 (8.9)	30/354 (8.5)	40/354 (11.3)

<sup>1</sup> IFA= Iron-Folic Acid group; MMN=Multiple Micronutrients group. LNS= Small Quantity Lipid-based Nutrient Supplement group. HFIA is Household Food Insecurity Access Score. N=total number of participants in the group in question; n = number of participants positive on the variable in question; % = percent of participants positive on the variable in question. Values are Mean ± SD [N] or n/N (%).

<sup>2</sup> Proxy indicators for household socioeconomic status; higher values represent higher socioeconomic status.

<sup>3</sup> HFIA (Household food insecurity access) is a proxy indicator for household food insecurity (Coates et al., 2007); higher values represent higher food insecurity

<sup>4</sup> Rapid Diagnostic Test (Clearview Malarial Combo, Vision Biotech, South Africa), which detected *P. falciparum* and non-*P. falciparum* histidine-rich protein-2

**TABLE 2**

Unadjusted hemoglobin, and iron status and inflammatory markers of pregnant Ghanaian women at baseline and 36 gestational weeks, by intervention group, and pair-wise comparison of groups<sup>1</sup>

Variable	IFA <sup>2</sup> [N=349]	MMN <sup>2</sup> [N=354]	LNS <sup>2</sup> [N=354]	P <sup>3</sup>	Comparison of IFA and MMN		Comparison of IFA and LNS		Comparison of MMN and LNS	
					Mean difference (95 % CI)	p	Mean difference (95 % CI)	p	Mean difference (95 % CI)	p
Hb, g/L										
Baseline	112 ± 13 [349]	111 ± 12 [354]	112 ± 12 [354]		-1 (-4, 1)		0 (-2, 2)		1 (-1, 3)	
36 GW	120 ± 11 [270]	117 ± 12 [291]	115 ± 12 [266]	<0.001	-3 (-6, -1)	0.002	-5 (-7, 3)	<0.001	-2 (-4, 1)	0.18
ZPP, µmol/mol heme										
Baseline	43 ± 28 [347]	46 ± 36 [354]	45 ± 33 [354]		3 (-3, 8)		2 (-4, 8)		-1 (-6, 5)	
36 GW	42 ± 30 [267]	49 ± 30 [291]	50 ± 29 [264]	<0.001	6 (0, 12)	<0.001	7 (1, 13)	<0.001	1 (-5, 7)	0.91
TfR, mg/L										
Baseline	4.0 ± 1.9 [338]	4.0 ± 1.7 [348]	4.3 ± 3.5 [346]		0.0 (-0.4, 0.5)		0.3 (-0.1, 0.8)		0.3 (-0.1, 0.8)	
36 GW	4.0 ± 1.3 [266]	4.6 ± 1.7 [291]	4.9 ± 1.8 [265]	<0.001	0.5 ( 0.2, 0.9)	<0.001	0.8 ( 0.5, 1.2)	<0.001	0.3 (-0.0, 0.6)	0.07
CRP, mg/L										
Baseline	7.9 ± 14 [338]	5.8 ± 8.3 [348]	6.9 ± 11 [346]		-2.1 (-4.1, -0.1)		-1.1 (-3.1, 1.0)		1.0 (-1.0, 3.0)	
36 GW	5.6 ± 18 [266]	5.7 ± 14 [291]	5.9 ± 18 [265]	0.85	0.1 (-3.3, 3.5)	0.95	0.3 (-3.2, 3.8)	0.97	0.2 (-3.2, 3.6)	0.84
AGP, g/L										
Baseline	0.7 ± 0.2 [338]	0.6 ± 0.2 [348]	0.6 ± 0.2 [346]		0.0 (-0.1, 0.0)		0.0 (-0.1, 0.0)		0.0 (0.0, 0.0)	

36 GW	0.5 ± 0.2 [266]	0.5 ± 0.2 [291]	0.5 ± 0.2 [265]	0.30	0.0 (0.0, 0.1)	0.65	0.0 (0.0, 0.0)	0.80	0.0 (-0.1, 0.0)	0.27
-------	--------------------	--------------------	--------------------	------	----------------	------	----------------	------	-----------------	------

<sup>1</sup>IFA= Iron-Folic Acid group; MMN=Multiple Micronutrients group; LNS= Small Quantity Lipid-based Nutrient Supplement group. AGP, CRP, GW, Hb, TfR, and ZPP are alpha-1 acid glycoprotein, C-reactive protein, gestational weeks, hemoglobin, transferrin receptor, and zinc protoporphyrin, respectively. Analyses are based on ANOVA (SAS, PROC GLM). Group means ( $\pm$ SD) and pair-wise mean difference (95% CI) were calculated using untransformed data. Except for Hb, all p-values for group or pair-wise comparisons were generated from log-transformed data; untransformed data were used for comparisons of mean Hb values. N=total number of participants in the group in question.

<sup>2</sup> Values are Mean  $\pm$  SD [N].

<sup>3</sup>P-values, with Tukey-Kramer adjustments, compare all three groups.

Number (percentage) of pregnant Ghanaian women with abnormal hemoglobin, and iron status and inflammatory markers at baseline and 36 weeks of gestation, by intervention group and pairwise relative risks (RR) between groups<sup>1</sup>

	IFA <sup>2</sup> [N=349]	MMN <sup>2</sup> [N=354]	LNS <sup>2</sup> [N=354]	P <sup>3</sup>	Comparison of IFA and MMN <sup>4</sup>		Comparison of IFA and LNS <sup>4</sup>		Comparison of MMN and LNS <sup>4</sup>	
					RR (95 % CI)	p	RR (95 % CI)	p	RR (95 % CI)	p
<b>Low Hb (&lt; 100 g/L)<sup>5</sup></b>										
Baseline	39/349 (11.2)	51/354 (14.4)	47/354 (13.3)		1.29 (0.81, 2.05)		1.19 (0.74, 1.91)		0.92 (0.59, 1.43)	
36 GW	6/270 (2.2)	17/291 (5.8)	21/266 (7.9)	0.019	2.63 (0.88, 7.86)	0.09	3.55 (1.22, 10.3)	0.013	1.35 (0.65, 2.83)	0.60
<b>Low Hb (&lt; 110 g/L)<sup>6</sup></b>										
Baseline	139/349 (39.8)	157/354 (44.4)	134/354 (37.9)		1.11 (0.90, 1.37)		0.95 (0.76, 1.19)		0.85 (0.69, 1.06)	
36 GW	38/270 (14.1)	69/291 (23.7)	88/266 (33.1)	<0.001	1.68 (1.10, 2.59)	0.011	2.35 1.56, 3.53)	<0.001	1.40 (1.01, 1.92)	0.038
<b>High Hb<sup>7</sup></b>										
Baseline	21/349 (6.0)	15/354 (4.2)	16/354 (4.5)		0.70 (0.33, 1.52)		0.75 (0.35, 1.60)		1.07 (0.47, 2.43)	
36 GW	42/270 (15.6)	32/291 (11.0)	28/266 (10.5)	0.15	0.71 (0.42, 1.18)	0.25	0.68 (0.40, 1.16)	0.20	0.96 (0.54, 1.70)	0.98
<b>Elevated ZPP<sup>8</sup></b>										
Baseline	40/347 (11.5)	54/354 (15.3)	46/354 (13.0)		1.32 (0.84, 2.09)		1.13 (0.70, 1.81)		0.85 (0.55, 1.32)	
36 GW	25/267 (9.4)	56/291 (19.2)	49/264 (18.6)	0.003	2.06 (1.21, 3.48)	0.003	1.98 (1.16, 3.40)	0.007	0.96 (0.64, 1.46)	0.98
<b>Elevated TfR<sup>9</sup></b>										
Baseline	29/338 (8.6)	29/348 (8.3)	37/346 (10.7)		0.97 (0.54, 1.75)		1.25 (0.72, 2.17)		1.28 (0.74, 2.23)	
36 GW	24/266 (9.0)	44/291 (15.1)	51/265 (19.2)	0.004	1.68 (0.96, 2.94)	0.08	2.13 (1.24, 3.67)	0.003	1.27 (0.82, 1.97)	0.40
<b>IDA<sup>10</sup></b>										
Baseline	20/338 (5.9)	22/348 (6.3)	22/346 (6.4)		1.07 (0.53, 2.15)		1.07 (0.53, 2.17)		1.01 (0.51, 1.99)	
36 GW	2/264 (0.8)	11/291 (3.8)	11/263 (4.2)	0.07	4.99 (0.83, 29.9)	0.09	5.52 (0.92, 33.1)	0.06	1.11 (0.42, 2.95)	0.97
<b>IDA<sup>11</sup></b>										
Baseline	36/338 (10.7)	37/348 (10.6)	37/346 (10.7)		1.00 (0.59, 1.68)		1.00 (0.60, 1.69)		1.01 (0.60, 1.68)	
36 GW	12/264 (4.5)	27/291 (9.3)	41/263 (15.6)	<0.001	2.04 (0.93, 4.49)	0.08	3.43 (1.63, 7.20)	<0.001	1.68 (0.97, 2.90)	0.07
<b>Elevated CRP only<sup>12</sup></b>										

Baseline	113/338 (33.4)	108/348 (31.0)	121/346 (35.0)		0.93 (0.72, 1.20)		1.05 (0.82, 1.34)		1.13 (0.87, 1.45)	
36 GW	49/266 (18.4)	70/291 (24.1)	59/265 (22.3)	0.26	1.31 (0.89, 1.93)	0.24	1.21 (0.81, 1.81)	0.52	0.93 (0.64, 1.33)	0.87
Elevated CRP+AGP <sup>13</sup>										
Baseline	26/338 (7.7)	15/348 (4.3)	11/346 (3.2)		0.56 (0.27, 1.17)		0.41 (0.18, 0.94)		0.74 (0.30, 1.84)	
36 GW	7/266 (2.6)	9/291 (3.1)	5/265 (1.9)	0.67	1.18 (0.37, 3.77)	0.94	0.72 (0.18, 2.79)	0.83	0.61 (0.17, 2.22)	0.64
Elevated AGP only <sup>14</sup>										
Baseline	3/338 (0.9)	5/348 (1.4)	5/346 (1.4)		1.62 (0.30, 8.88)		1.63 (0.30, 8.93)		1.01 (0.23, 4.38)	
36 GW	1/266 (0.4)	0/291 (0.0)	1/265 (0.4)	1.00						

<sup>1</sup>IFA= Iron-Folic Acid group; MMN=Multiple Micronutrients group; LNS=Small Quantity Lipid-based Nutrient Supplement group. AGP, CRP, GW, Hb, IDA, TfR, and ZPP are alpha-1 acid glycoprotein, C-reactive protein, gestational weeks, hemoglobin, iron deficiency anemia, transferrin receptor, and zinc protoporphyrin, respectively. N=total number of participants in the group in question.

<sup>2</sup>Values are number of participants positive on the variable in question/N (% of participants positive on the variable in question).

<sup>3</sup>P-values compare all three groups, with Tukey-Kramer adjustment, using logistic regression (SAS PROC LOGISTIC).

<sup>4</sup>Relative Risks, RR(95% CI) and their p-values are based on Poisson regression (Spiegelman and Hertzmark, 2005).

<sup>5</sup>Nestel and INACG Steering Committee, 2002, WHO, 2007, WHO/UNICEF/UNU, 2001.

<sup>6</sup>WHO, 2011.

<sup>7</sup>Hb >130 g/L (Pena-Rosas et al., 2012).

<sup>8</sup>ZPP > 60.0 (μmol/mol heme).

<sup>9</sup>TfR >6.0 mg/L (Pfeiffer et al., 2007, Vandevijvere et al., 2013).

<sup>10</sup>Hb <100 g/L (Nestel and INACG Steering Committee, 2002, WHO, 2007, WHO/UNICEF/UNU, 2001) and at least one marker of iron deficiency (ZPP > 60.0 (μmol/mol heme) or TfR >6.0 mg/L (Pfeiffer et al., 2007, Vandevijvere et al., 2013)).



<sup>11</sup>Hb <110 g/L (WHO, 2011) and at least one marker of iron deficiency (ZPP > 60.0 (μmol/mol heme) or TfR >6.0 mg/L (Pfeiffer et al., 2007, Vandevijvere et al., 2013)).

<sup>12</sup>CRP >5.0 mg/L and AGP not >1.0 g/L (Thurnham and McCabe).

<sup>13</sup> CRP >5.0 mg/L and AGP >1.0 g/L (Thurnham and McCabe).

<sup>14</sup> CRP not > 5.0 mg/L and AGP >1.0 g/L (Thurnham and McCabe).

**TABLE 4**Effect of intervention on iron status and inflammatory outcomes of pregnant Ghanaian women, stratified by baseline characteristics<sup>1</sup>

Outcomes	IFA <sup>2</sup>	MMN <sup>2</sup>	LNS <sup>2</sup>	P <sup>3</sup>	P <sup>4</sup>	Comparison of MMN and IFA (n = 349)		Comparison of LNS and IFA (n = 354)		Comparison of LNS and MMN (n = 354)	
						Difference or RR	P	Difference or RR	P	Difference or RR	P
ZPP, $\mu\text{mol/mol heme}$ <sup>5</sup>											
Baseline anemia				0.099							
No <sup>†</sup>	44 $\pm$ 2 [233]	47 $\pm$ 2 [245]	48 $\pm$ 2 [226]		0.09	3 (-2, 9)	0.34	5 (-0, 11)	0.08	2 (-4, 8)	0.71
Yes	36 $\pm$ 6 [26]	54 $\pm$ 5 [41]	53 $\pm$ 5 [33]		0.013	18 (3, 34)	0.017	18 (1, 34)	0.032	-1 (-16, 14)	0.99
Elevated baseline ZPP				0.013							
No	42 $\pm$ 2 [230]	46 $\pm$ 2 [242]	46 $\pm$ 2 [226]		0.19	3 (-2, 9)	0.34	4 (-2, 10)	0.20	1 (-5, 7)	0.94
Yes	47 $\pm$ 5 [29]	64 $\pm$ 5 [44]	72 $\pm$ 5 [33]		0.001	18 (3, 32)	0.015	25 (9, 41)	0.001	8 (-7, 22)	0.43
Elevated baseline TfR				0.090							
No	43 $\pm$ 2 [239]	47 $\pm$ 2 [261]	49 $\pm$ 2 [233]		0.06	4 (-2, 9)	0.28	6 (0, 12)	0.048	2 (-4, 8)	0.65
Yes	41 $\pm$ 6.5 [20]	63 $\pm$ 6 [25]	59 $\pm$ 6 [26]		0.017	22 (3, 41)	0.017	18 (-1, 36)	0.07	-4 (-22, 14)	0.84
GA at enrolment				0.041							
At 10 <sup>th</sup> percentile <sup>‡</sup>	46 $\pm$ 3	51 $\pm$ 3	44 $\pm$ 3		0.28	5 (-5, 15)	0.44	-1 (-11, 9)	0.97	-6 (-16, 4)	0.29
At 90 <sup>th</sup> percentile	40 $\pm$ 3	48 $\pm$ 3	54 $\pm$ 3		0.002	7 (-1, 16)	0.12	13 (5, 22)	0.001	6 (-3, 15)	0.25
Assets score				0.061							
At 10 <sup>th</sup> percentile	40 $\pm$ 3	46 $\pm$ 3	54 $\pm$ 3		0.003	5 (-4, 15)	0.38	14 (4, 24)	0.002	8 (-1, 18)	0.11
At 90 <sup>th</sup> percentile	46 $\pm$ 3	50 $\pm$ 3	46 $\pm$ 3		0.37	5 (-4, 13)	0.44	-0 (-9, 9)	1.00	-5 (-14, 4)	0.46
Elevated ZPP <sup>6</sup>											
GA at enrolment				0.057							
At 10 <sup>th</sup> percentile <sup>§</sup>	11.1 (5.7, 20.5)	23.2 (15.3, 33.6)	11.6 (6.4, 20.0)		0.06	3.0 (0.9, 4.6)	0.12	1.0 (0.4, 2.5)	1.00	0.5 (0.2, 1.1)	0.11
At 90 <sup>th</sup> percentile	5.8 (2.9, 11.4)	14.9 (9.6, 22.3)	20.7 (14.1, 29.2)		0.004	2.4 (1.0, 5.7)	0.049	3.4 (1.5, 7.7)	0.003	1.4 (0.8, 2.5)	0.46

Outcomes	IFA <sup>2</sup>	MMN <sup>2</sup>	LNS <sup>2</sup>	P <sup>3</sup>	P <sup>4</sup>	Comparison of MMN and IFA (n = 349)		Comparison of LNS and IFA (n = 354)		Comparison of LNS and MMN (n = 354)	
						Difference or RR	P	Difference or RR	P	Difference or RR	P
Elevated baseline TfR				0.049							
No <sup>¶</sup>	17/234 (7.4)	44/256 (17.4)	32/229 (14.0)		0.008	2.2 (1.2, 4.0)	0.006	1.8 (1.0, 3.4)	0.08	0.8 (0.5, 1.3)	0.61
Yes	1/20 (3.0)	3/25 (10.0)	8/25 (31.3)		0.014	1.3 (0.4, 4.6)	0.41	2.7 (0.8, 9.4)	0.016	2.1 (0.7, 6.1)	0.14
TfR, mg/L <sup>7</sup>											
Baseline anemia				0.079							
No	4.1 ± 0.1 [232]	4.5 ± 0.1 [245]	4.8 ± 0.1 [226]		<0.001	0.4 (0.1, 0.8)	0.004	0.7 (0.4, 1.0)	<0.001	0.3 (-0.0, 0.6)	0.11
Yes	3.5 ± 0.3 [26]	4.8 ± 0.3 [41]	5.0 ± 0.3 [33]		<0.001	1.3 (0.4, 2.1)	0.002	1.5 (0.6, 2.4)	<0.001	0.2 (-0.6, 1.1)	0.76
Elevated baseline AGP				0.051							
No	4.1 ± 0.1 [238]	4.6 ± 0.1 [272]	4.9 ± 0.1 [250]		<0.001	0.6 (0.3, 0.9)	<0.001	0.8 (0.5, 1.2)	<0.001	0.3 (-0.02, 0.6)	0.07
Yes	4.0 ± 0.4 [20]	4.0 ± 0.4 [14]	3.3 ± 0.5 [9]		0.503	-0.0 (-1.2, 1.2)	1.00	-0.7 (-2.1, 0.8)	0.51	-0.66 (-2.2, 0.8)	0.56
GA at enrolment				0.016							
At 10 <sup>th</sup> percentile	4.4 ± 0.2	5.0 ± 0.2	4.8 ± 0.2		0.023	0.9 (0.1, 1.1)	0.019	0.4 (-0.1, 0.9)	0.16	-0.2 (-0.7, 0.3)	0.64
At 90 <sup>th</sup> percentile	3.8 ± 0.1	4.3 ± 0.1	4.9 ± 0.1		<0.001	0.5 (0.0, 0.9)	0.041	1.1 (0.7, 1.6)	<0.001	0.7 (0.2, 1.1)	0.002
Elevated TfR <sup>8</sup>											
HFIA score				0.047							
At 10 <sup>th</sup> percentile	5.3 (2.9, 9.7)	15.7 (10.8, 22.2)	15.3 (10.5, 21.7)		0.004	2.3 (1.1, 4.7)	0.006	2.5 (1.2, 5.0)	0.007	1.1 (0.6, 1.8)	1.00
At 90 <sup>th</sup> percentile	9.7 (5.1, 17.5)	4.9 (1.7, 13.5)	18.7 (10.3, 31.5)		0.06	0.7 (0.2, 2.0)	0.50	1.9 (0.8, 4.6)	0.27	2.8 (0.9, 8.7)	0.06
Elevated CRP <sup>9</sup>											
Elevated baseline ZPP				0.008							
No	41/225 (18.3)	72/238 (30.1)	47/221 (21.4)		0.013	1.6 (1.1, 2.4)	0.014	1.2 (0.8, 1.8)	0.71	0.7 (0.5, 1.1)	0.11
Yes	9/29 (30.1)	6/43 (13.5)	11/33 (32.0)		0.09	0.5 (0.2, 1.1)	0.17	1.1 (0.5, 2.1)	0.99	2.1 (0.9, 4.9)	0.10

Outcomes	IFA <sup>2</sup>	MMN <sup>2</sup>	LNS <sup>2</sup>	P <sup>3</sup>	P <sup>4</sup>	Comparison of MMN and IFA (n = 349)		Comparison of LNS and IFA (n = 354)		Comparison of LNS and MMN (n = 354)	
						Difference or RR	P	Difference or RR	P	Difference or RR	P
Elevated AGP <sup>10</sup>											
Assets score				0.091							
At 10 <sup>th</sup> percentile	1.0 (0.2, 4.8)	1.0 (0.2, 4.5)	2.8 (0.9, 8.4)		0.39	1.0 (0.1, 9.8)	1.00	2.8 (0.6, 14.1)	0.51	2.8 (0.3, 28.0)	0.48
At 90 <sup>th</sup> percentile	4.3 (1.5, 11.3)	4.4 (1.7, 10.8)	0.7 (0.1, 3.9)		0.14	1.0 (0.2, 4.5)	1.00	0.16 (0.0, 1.3)	0.16	0.2 (0.0, 1.4)	0.14

<sup>1</sup>IFA= Iron-Folic Acid group; MMN=Multiple Micronutrients group; LNS=Small Quantity Lipid-based Nutrient Supplement group. GA, Hb, HFIA, TfR, and ZPP are gestation age, hemoglobin, household food insecurity access, transferrin receptor, and Zinc protoporphyrin, respectively. Baseline anemia is Hb <100 g/L (Nestel and INACG Steering Committee, 2002, WHO, 2007, WHO/UNICEF/UNU, 2001). Analyses are based on ANCOVA (SAS PROC MIXED, with SLICE option) for continuous outcomes or logistic regression (SAS PROC GLIMMIX, with SLICE option) for binary outcomes. Linear regression modeling was used to predict the outcome at the 10<sup>th</sup> and 90<sup>th</sup> percentile of continuous baseline effect modifiers.

<sup>2</sup>Values are mean  $\pm$  SE [total number of participants], or mean  $\pm$  SE, or percent of participants positive on the variable in question (95% CI) or number of participants positive on the variable in question/total number of participants (percent of participants positive on the variable in question). Values are adjusted for variables significantly associated with the outcome variable in bivariate analysis.

<sup>3</sup>P-values are for interaction with the outcome in question

<sup>4</sup>P-values compare all three groups in each stratum.

<sup>5</sup>ZPP values are adjusted for age, parity, season at enrolment, and baseline Hb, ZPP, and TfR concentrations.

<sup>6</sup>Elevated ZPP (ZPP >60  $\mu$ mol/mol heme) adjusted for age, education, and baseline Hb, ZPP and TfR concentrations.

<sup>7</sup>TfR values are adjusted for age, season at enrolment, and baseline Hb, ZPP, TfR and AGP concentrations.

<sup>8</sup>Elevated TfR (TfR > 6.0 mg/L) adjusted for age, gestational age enrolment, season at enrolment, and baseline Hb, ZPP, and TfR concentrations.

<sup>9</sup>Elevated CRP (CRP > 5.0 mg/L) adjusted for BMI, household food insecurity access score, and baseline ZPP, AGP and CRP concentrations.

<sup>10</sup>Elevated AGP (AGP > 1.0 g/L) adjusted for age, parity, and season as enrolment.

<sup>†</sup>Values are group mean  $\pm$  SE [number of participants], and difference in means (95% CI) and p-values. All such values.

<sup>‡</sup>Values are group mean  $\pm$  SE, and difference in means (95% CI) and p-values. All such values.

<sup>§</sup>Values are group % (95% CI), and relative risk (95% CI) and p-values. All such values.

<sup>¶</sup>Values are number of participant (%), and relative risk (95% CI) and p-values. All such values.